



MODERN IDEAS ABOUT THE DIAGNOSIS AND TREATMENT OF ENDOMETRIOSIS

Umidova Nigora Nabi qizi, Aminova Sevinch Narzulloyevna.

Bukhara State Medical Institute.

About the author: Umidova Nigora Nabi qizi – assistant department of histology, cytology, embryology.

Summary: Endometriosis is one of the most common gynecological diseases, which reflects the medical and social significance of the problem of effective diagnosis and treatment. To date, the causes, diagnosis and treatment of this disease remain the subject of controversy. The pathogenesis of the disease is multifactorial and has not been studied enough, non-invasive examination methods have a relative diagnostic value, so modern therapeutic approaches often do not provide a complete cure.

Keywords: endometriosis, diagnostics, proteins, genes, mikro RNK

СОВРЕМЕННЫЕ ПРЕДСТАВЛЕНИЯ О ДИАГНОСТИКЕ И ЛЕЧЕНИИ ЭНДОМЕТРИОЗА

Умидова Нигора Наби кизи, Аминова Севинч Нарзуллоевна.

Бухарского государственного медицинского института.

Об авторах: Умидова Нигора Наби кызы – ассистент кафедры гистологии, цитологии, эмбриологии.

Резюме: Эндометриоз является одним из наиболее распространенных гинекологических заболеваний, что отражает медицинскую и социальную значимость проблемы эффективной диагностики и лечения. На сегодняшний день причины возникновения, диагностика и лечение этого заболевания остаются предметом споров. Патогенез заболевания имеет многофакторный характер и исследован недостаточно, неинвазивные методы обследования



имеют относительную диагностическую ценность, поэтому современные терапевтические подходы часто не обеспечивают полного излечения.

Ключевые слова: эндометриоз, диагностика, белки, гены, микро РНК

Genital endometriosis remains one of the most urgent problems of modern gynecology. External genital endometriosis (EGE) occurs in 5-10% of women of reproductive age. The frequency of detection of this disease during laparoscopy in order to clarify the cause of infertility is 45-55%. According to modern concepts, this pathology is one of the main causes of female infertility. Demonstrative indicators of the significance of endometriosis for reproductive health include the fertility rate, which is calculated as the ratio of the number of births to the number of women of reproductive age. This coefficient in healthy women is 0.15–0.2, and in patients with endometriosis it is 0.02–0.10. However, the issue of managing patients with infertility after surgical treatment remains relevant. R. Maheux and co-authors believe that the problem of endometriosis is one of the central problems in gynecology, along with inflammatory diseases of the pelvic organs and uterine myoma (Baskakov V.P. et al.; Adamyan L.V. et al.). The frequency of endometriosis does not tend to decrease and currently reaches 10% in the population (Guidice L.N.).

The progressive course of the disease leads to the occurrence of pain, menstrual disorders, infertility, psycho-emotional changes, and, as a result, a decrease in the quality of life of patients (Kira E. F., Ryabtseva M. V.).

The problem of early diagnosis of various forms of endometriosis is currently particularly acute, since there is still no reliable non-invasive diagnostic method or any pathognomonic clinical tests that could make an accurate diagnosis and identify the localization of endometriosis in the early stages of the disease.

External genital endometriosis (EGE) occurs in 5-10% of women of reproductive age. The frequency of detection of this disease during laparoscopy in order to clarify the cause of infertility is 45-55%. According to modern concepts, this pathology is one of the main causes of female infertility. In Russia, endometriosis among the causes of infertility ranks second after tubal-peritoneal infertility.



Demonstrative indicators of the significance of endometriosis for reproductive health include the fertility rate, which is calculated as the ratio of the number of births to the number of women of reproductive age.

This coefficient in healthy women is 0.15–0.2, and in patients with endometriosis it is 0.02–0.10. Recently, there has been a tendency to increase the frequency of severe forms of endometriosis. The main complaints of patients with severe forms of endometriosis are chronic pelvic pain, leading to a decrease in the working capacity and quality of life of women, and infertility, when young women cannot perform their reproductive function. This problem has a negative impact in the social, demographic and personal spheres.

According to Research Foundation, up to 17% of women in the study population reported having or suspected endometriosis. The average age of patients at diagnosis is from 28 to 35 years, while the average duration from the onset of symptoms to the diagnosis of endometriosis is about 6 years, and according to some sources - from 7 to 10 and even up to 12 years.

The study of the proteome may revolutionize the understanding of the etiopathogenesis of this disease and its localizations. Since endometriosis is an elusive disease, it is extremely important to identify high-precision markers for the timely diagnosis of this pathological process.

Given the steady increase in the incidence of endometriosis, which leads to a decrease in the quality of life, and sometimes to disability in women of predominantly reproductive age, the lack of reliable clinical criteria and specific, high-precision markers of the disease that can detect endometriosis in the early stages, it is necessary to create new diagnostic criteria and a prognostic model with the purpose of timely diagnosis of endometriosis and its localization.

Endometriosis is a chronic gynecological disease, representing a benign growth outside the uterine cavity of tissue similar in morphological and functional properties to the endometrium. The main clinical manifestations of the disease are persistent pain and infertility. Pain, as one of the symptoms of endometriosis, disrupts not only the physical and psychological state of a woman, but ultimately disrupts her



social and reproductive well-being. It should be noted that heterotopias are only similar to endometrial tissue, since the molecular genetic defects that characterize them contribute to infiltrative growth with penetration into surrounding tissues.

Reduced apoptosis and the absence of a connective tissue capsule, as well as the ability to metastasize, make it possible to compare it with a tumor process.

Another important and, at present, generally recognized fact, which has been established both in an experimental model of endometriosis in mice and in the abdominal cavity of patients with endometriosis, can be considered the development of the so-called local aseptic inflammation and dysfunction of immunocompetent cells. Damage to biological molecules (lipids, cell membranes, proteins, etc.) by highly reactive oxygen compounds is the basis for the pathogenesis of many diseases, and endometriosis is no exception.

The similarity of molecular biological indicators of neoangiogenesis, apoptosis, proliferation, invasion and expression of growth factors in areas of endometriosis and autologous hyperplastic endometrium has been proven. Peculiarities of expression of matrix metalloproteinases and their inhibitor in stromal and parenchymal elements were revealed depending on the clinical activity of adenomyosis and recurrence of endometrioid cysts. It has been established that the clinical activity of endometriosis is due to the activity of molecular biological processes involved in the genesis of the disease.

It has been determined that the stromal component in clinically active adenomyosis prevails over the epithelial component not only in quantitative terms, but also in terms of the severity of molecular biological indicators of proliferation, neoangiogenesis, invasion and expression of growth factors, which reflects the degree of maturity of the stroma, indicates remodeling and its stimulating role in the development of internal endometriosis, i.e., confirms the most important, aggressive role of the stroma in the development of adenomyosis.

Разработан способ прогнозирования клинической активности аденомиоза, основанный на выявлении иммуногистохимических показателей апоптоза, инвазии, неоангиогенеза, пролиферации и экспрессии факторов



роста в соскобе эндометрия (с учетом выявленной общности молекулярных показателей в ткани аденомиоза и аутологичном гиперплазированном эндометрии).

ЛИТЕРАТУРА / REFERENCES:

1. Саркисова, Л. В., Каюмова, Г. М., & Умидова, Н. Н. (2018). Морфологические изменения фетоплацентарного комплекса при герпетической инфекции. *Тиббиётда янги кун*, 188-191.
2. Каюмова, Г. М., Саркисова, Л. В., & Умидова, Н. Н. (2018). Современные взгляды на проблему преждевременных родов. *Тиббиётда янги кун*, 183-185.
3. Саркисова, Л. В., & Умидова, Н. Н. (2018). Premature birth in the modern aspect. *Новый день в медицине*, 3, 23.
4. Toyqulovna, K. M. T. K. M., & Nabievna, U. N. (2023). THE ROLE OF GENETIC DETERMINANTS IN THE OCCURRENCE OF HYPERPLASTIC PROCESSES OF THE REPRODUCTIVE SYSTEM OF WOMEN'S MENOPAUSAL AGE. *Journal of Advanced Zoology*, 44(S2), 3724-3730.
5. Саркисова, Л. В., Умидова, Н. Н., Муаззамов, Б. Б., Муаззамов, Б. Р., & Ахророва, Л. Б. (2019). Пути улучшения способов профилактики и лечения анемии беременных. *Новый день в медицине*, (2), 275-279.
6. Baxtiyorovna, N. N. (2023). Modern Aspects of Early Diagnosis and Effectiveness of Treatment of Endometriosis. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 23-28.
7. Baxtiyorovna, N. N. (2023). Modern View on the Diagnosis and Treatment of Endometriosis. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 20-22.
8. Hamdamova, M. T., & Umidova, N. N. (2023, September). THE ROLE OF APOPTOSIS MARKERS AND ANGIOGENESIS REGULATORS IN THE



PATHOGENESIS OF GENITAL ENDOMETRIOSIS. In *International Conference on Medicine and Life Sciences* (pp. 47-48).

9. Khamdamova, M. T., & Umidova, N. N. (2023, September). MEDICAL AND SOCIAL ASPECTS OF GENITAL, ENDOMETRIOSIS. In *International Conference on Medicine and Life Sciences* (pp. 49-50).

10. Умидова, Н. Н. (2023). Медицинские И Социальные Аспекты Генитального, Эндометриоза. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(5), 416-418.

11. Samandarovna, S. Z., & Hikmatovna, A. M. (2023). Treatment Efficiency Analysis Pregnant, With Anemia of Varying Severity. *Scholastic: Journal of Natural and Medical Education*, 2(2), 214-218.

12. Uchqunovna, N. M., & Ixtiyarovna, N. O. (2022). Treatment of Pregnant People with Various Anemia Disease at Random in Humans. *Central Asian Journal of Literature, Philosophy and Culture*, 3(12), 153-159.

13. Саркисова, Л., & Умидова, Н. (2019). Анализ эффективности лечения беременных, с анемией различной степени тяжести. *Журнал вестник врача*, 1(4), 115-118.

14. Umidova, N. N. (2019). THE ROLE OF INDICATORS OF FERROKINETICS AND ENDOGENOUS ERYTHROPOIETIN IN ANEMIA OF PREGNANT WOMEN. *Journal of Asian Medical Student Association*, 7(2), 48-50.

15. Sarkisova, L. V., Umidova, N. N., & Ro'ziyeva, D. U. (2019). Treatment efficiency analysis pregnant, with anemia of varying severity. *Новый день в медицине*, (4), 290-294.